Table 1: Primary and secondary endpoints from the ROAR study

	ITT ATC cohort (n=36)		BRAF V600E assessable (n=33)	
	Investigator assessed	Independent assessment	Investigator assessed	Independent assessment
ORR	56%	53%	61%	58%
(95% CI)	(38.1% to	(35.5% to	(42.1% to	(39.2% to
N N	72.1%)	69.6%)	77.1%)	74.5%)
CR	8% (3)	6% (2)	9% (3)	6% (2)
(patients)		656 WKD8. 65 WKP. 19	apressed some.	Andrew Constant
PR	47% (17)	47% (17)	52% (17)	52% (17)
(patients)	No 020	16 15to	N 155	N 850
Median	14.4 months	13.6 months	NR	NR
DOR	(7.4 to 43.6	(3.8 to not		
(95% CI)	months)	reached)		
Median	6.7 months	5.5 months	NR	NR
PFS	(4.7 to 13.8	(3.7 to 12.9		
(95% CI)	months)	months)		
Overall	14.5 months	NR	NR	NR
survival	(95% CI 6.8 to	(Value of the Control	300,000	V-10000
	23.2 months)			
ATC: anapla	astic thyroid cance	er; CI: confidence	interval; CR: com	plete response;
		uration of respons		
		nse rate; PFS: pro		