



Epoetin zeta (Retacrit®)

For the reduction of exposure to allogeneic blood transfusions in adult non-iron deficient patients prior to major elective orthopaedic surgery, having a high perceived risk for transfusion complications. Use should be restricted to patients with moderate anaemia (e.g. Hb 10-13 g/dl) who do not have an autologous predonation programme available and with expected moderate blood loss (900 to 1800 ml)

Hospira UK Ltd

November 2011

Statement of Advice

ADVICE: in the absence of a submission from the holder of the marketing authorisation, epoetin zeta (Retacrit®) cannot be endorsed for use within NHS Wales for the reduction of exposure to allogeneic blood transfusions in adult non-iron deficient patients prior to major elective orthopaedic surgery, having a high perceived risk for transfusion complications

Advice context:

The All Wales Medicines Strategy Group (AWMSG) takes into account the National Institute for Health and Clinical Excellence (NICE) future work programme when considering whether a product will be appraised. To avoid duplication of effort, AWMSG would not normally consider undertaking an appraisal if NICE intend to publish final advice for the same product within twelve months of the projected Form B submission date. AWMSG advice is interim to that of NICE, should NICE subsequently publish guidance.

The above medicine cannot be endorsed for use within NHS Wales as an appraisal by NICE or AWMSG has not been undertaken. The medicine should NOT be prescribed routinely within NHS Wales for the indication stated above.

In the absence of guidance issued by NICE or AWMSG, clinicians should continue to exercise their clinical judgement when providing care for an individual patient. This should be in consultation with the patient and/or guardian or carer, based on the best available evidence.

This statement will be removed on receipt of a submission (i.e. the full submission [Forms A and B] or limited submission [Forms A and C]) or when final NICE guidance becomes available.